2011 Recommended Immunizations for Children from Birth Through 6 Years Old

Birth	1 month	2 months	4 months	6 months	12 months	15 months	18 months	19–23 months	2–3 years	4–6 years
HepB	HepB			HepB						
		RV	RV	RV						
		DTaP	DTaP	DTaP		DTaP				DTaP
		Hib	Hib	Hib	Hib					
		PCV	PCV	PCV	PCV§					
		IPV	IPV	IPV						IPV
				Influenza (Yearly)*						
					MMR					MMR
					Varicella					Varicella
					HepA, 2 doses§					

Shaded boxes indicate the vaccine can be given during shown age range.

NOTE: If your child misses a shot, you don't need to start over, just go back to your child's doctor for the next shot. The doctor will keep your child up-to-date on vaccinations. Talk with your doctor if you have questions.

FOOTNOTES

§ HepA vaccination is recommended for high-risk children older than 2 years. Children with certain medical conditions may also need a dose of meningococcal vaccine (MCV4) and pneumococcal vaccine (PPSV). HepA vaccination may be administered to any child older than 2 years for whom immunity is desired. See vaccine-specific recommendations at http://www.cdc.gov/vaccines/pubs/ACIP-list.htm.

* Two doses given at least four weeks apart are recommended for children aged 6 months through 8 years of age who are getting a flu vaccine for the first time. Children who only got one dose in their first year of vaccination should get two doses the following year.

See back page for more information on vaccine-preventable diseases and the vaccines that prevent them.

For more information, call toll free
1-800-CDC-INFO (1-800-232-4636)
or visit
http://www.cdc.gov/vaccines

U.S. Department of Health and Human Services
Centers for Disease Control and Prevention

AMERICAN ACADEMY OF FAMILY PHYSICIANS
STRONG MEDICINE FOR AMERICA

American Academy of Pediatrics
DEDICATED TO THE HEALTH OF ALL CHILDREN™